


Eur J Vasc Endovasc Surg 21, 51–56 (2001)

doi:10.1053/ejvs.2000.1262, available online at <http://www.idealibrary.com> on 

## Smoking, but not Lipids, Lipoprotein (a) and Antibodies Against Oxidised LDL, is Correlated to the Expansion of Abdominal Aortic Aneurysms

J. S. Lindholt<sup>\*1</sup>, N. H. H. Heegaard<sup>2</sup>, S. Vammen<sup>1</sup>, H. Fasting<sup>1</sup>, E. W. Henneberg<sup>1</sup> and L. Heickendorff<sup>3</sup>

<sup>1</sup>Department of Vascular Surgery, Hospital of Viborg, Viborg, Denmark, <sup>2</sup>Department of Autoimmunology, Statens Serum Institute, Copenhagen, Denmark, <sup>3</sup>Department of Clinical Biochemistry, Aarhus Amtssygehus, University of Aarhus, Denmark

**Objectives:** to study the role of smoking, lipids, lipoprotein (a), and autoantibodies against oxidised low density lipoprotein (Ab-oxLDL) in the expansion of small abdominal aortic aneurysms (AAA). To study the role of Ab-oxLDL and Lp(a) in the progression of lower limb atherosclerosis.

**Methods and Materials:** one hundred and thirty-eight male patients with AAA were interviewed, examined, and their serum lipids and S-Ab-oxLDL determined. Of these, 117 were followed annually with ultrasound and underwent control scans and blood pressure measurements for a mean of 2.5 (range 1–5) years.

**Results:** initial AAA size, smoking and level of triglycerides were positively correlated to increased aneurysmal expansion, while beta-blocker medication was associated with decreased expansion. Besides initial AAA size, only smoking had persisting significance after adjustment of the other significant variables. Initial ankle brachial pressure index (ABI) and Lp(A) but not Ab-oxLDL were significantly correlated to ABI change.

**Conclusion:** smoking cessation may inhibit aneurysmal expansion. Lipids seem to play a minor role in the progression of AAA.

**Key Words:** Smoking; Lipids; Oxidized low density lipoprotein; Lipoprotein (a); Abdominal aortic aneurysm; Atherosclerosis; Progression.

### Introduction

Predicting the expansion of small abdominal aortic aneurysms (AAA) is important for clinical management. Lipoprotein (A) (LpA) levels are related to atherosclerotic disease but the relationship between LpA-levels and the progression of AAA has not been analysed.<sup>1</sup>

Oxidized LDL is implicated in the initial damage of the vascular endothelium.<sup>2–4</sup> The concentration of antibodies against oxidised LDL (Ab-oxLDL) is elevated in patients with acute myocardial infarction (AMI)<sup>2</sup> and carotid stenosis.<sup>2,3</sup> The correlation was increased in cases with hypertriglyceridaemia.<sup>5</sup> The role of oxidised LDL in the pathogenesis of AAA is unknown.

The aim of the study was to relate AAA expansion with lipids, Lp(A) and Ab-oxLDL levels.

### Material and Methods

Details of our AAA screening programme have been published previously.<sup>6</sup>

Patient sera were analysed for IgG antibodies against oxidised LDL by means of an enzyme-linked immunosorbent assay (ELISA) using oxLDL-coated microtitre plates, as previously described.<sup>7,8</sup>

P-Lipoprotein a (Lp(a)) was quantified with earlier reported and validated immunoradiometric assay (IRMA) (Pharmacia Diagnostics, Sweden).<sup>9</sup>

S-total-cholesterol, S-high density lipoprotein (HDL), and S-triglycerides were determined by commercial assays, while S-LDL was calculated as: S-cholesterol – S-HDL + (0.45\*S-triglycerides).<sup>10</sup>

One hundred and seventy men had an AAA diagnosed at screening (4.0%).<sup>11,12</sup> Complete data were available in 138 cases (82%). Twenty-one were above 5 cm and referred for surgery, and the remaining 87% were offered annual ultrasound scans. Of these, 124 have now been followed for 1–5 years (average 2.5 years). Blood samples were complete in 117 (94%) of these patients. The mean expansion rate was mean 2.6 mm/year (SD:2.6). The ABI change was on average –2.2%/year (SD:6.8). The interobserver variations (def.: 2SD) of aortic diameter and ABI-change were 1.68 mm,<sup>13</sup> 0.16,<sup>14</sup> respectively. The intra-assay coefficients of variation of Ab-oxLDL and Lp(a) were 10% and 2.2%, respectively. The interassay coefficients

\* Please address all correspondence to: J. S. Lindholt, Lærkevej 11, 8900 Randers, Denmark.

of variation of Ab-oxLDL and Lp(a) were 25% and 3.1%, respectively.<sup>9</sup> The same assays were used for all the samples.

Also, Wilcoxon's rank sum test and Spearman's rank correlation analysis were used for the univariate analyses, while multiple linear regression analyses only were performed in significant univariate findings after securing that additivity, linearity, variance homogeneity and normality of residuals were reasonable fulfilled. Unfortunately, Lp(a) even after transformation failed to be reasonably normally distributed. The statistical software package used was SPSS 10.0.

## Results

Table 1 shows the relationship between measured biochemical and clinical variables and use of beta-blockers, diet and smoking.

The relationship between lipids, Lp(a), Ab-oxLDL, AAA expansion, and ABI are tested upon beta-blocker medication, dietary habits, supplements of vitamin and Q10 (antioxidant) tablets, and smoking habits. Beta-blockers were associated with an atherogenic lipid profile, lower expansion and better ABI. However, these relationships disappeared after adjustment for smoking. Smoking was associated with aneurysmal expansion but not with change in ABI.

Besides the initial AAA size, S-triglycerides were correlated to expansion, while only the initial ABI and lipoprotein (A) were associated with the ABI-change-rate (Table 2). The initial AAA size remained significantly correlated to expansion after adjustment for smoking, while the correlation between triglycerides and expansion disappeared after adjustment for smoking.

## Discussion

Only smoking and initial AAA size was correlated to expansion, while only initial ABI and concentration of lipoprotein (A) were correlated to ABI change rate.

Most reports studying the relation between aneurysmal size and expansion have also found initial AAA size of the AAA predictive for expansion<sup>15-19</sup> but we were not able to confirm earlier reports of increased expansion with increasing diastolic brachial blood pressure<sup>20,21</sup> or systolic ankle blood pressure.<sup>16</sup>

Smoking has frequently been reported to be associated with AAA in case control studies,<sup>22-28</sup> but

apparently has only been reported once<sup>29</sup> to be correlated with expansion. This may be due to the association with the frequently coexisting atherosclerotic diseases, or due to the use of unreliable data obtained by interviews. The latter bias could be omitted by cotinine measurements.<sup>23,29</sup> Our smoking data was obtained from a questionnaire that was completed at home before the initial consultation. This may have limited the frequency of invalid data. Furthermore, smoking data was obtained at the following annual control scans; only one of the "non-smokers" started smoking. In spite of recommendations to stop smoking, unfortunately only one patient did so. This consistency of data suggests reliable data concerning smoking habits.

Beta-blockers were analysed because of earlier reports of lower expansion among beta-blocker-users.<sup>21,30,31</sup> The possible effect of beta-blockage is hypothesised to be caused by increased cross-linkage of collagen- and elastin.<sup>32</sup> Our observations confirm these previous observations. However, adjustment for smoking removed the significant associations, indicating that the apparently benefits of beta-blockage could be, at least partly, due to limited smoking among beta blocker users. This seems logical because acute myocardial infarction or other cardiac events are one of the most common reasons for beta-blockage. Such events could easily be more motivating to stop smoking than general health campaigns or an accidental finding of a asymptomatic small AAA. Furthermore, our earlier experience with a randomised intervention study with propranolol versus placebo could not show any effect. However, this was mainly because of a high frequency of side effects and decreased quality of life among those receiving propranolol, resulting in a low patient compliance in this study.<sup>33</sup>

A significant correlation between s-triglycerides and aneurysmal expansion was noticed. However, the correlation disappeared after adjustment for any of the other significant variables; beta-blockage, smoking, or initial AAA size. In spite of many earlier prospective studies of small AAA, only one study apart from our own seems to have reported increased expansion associated to hyperlipidaemia.<sup>34</sup> Furthermore, the association seems controversial; some reports have observed increased levels of various lipids in AAA-patients<sup>22,35,36</sup> while others have failed to do so.<sup>23-25</sup> These different reports may be explained by different patient populations with different frequencies of coexisting atherosclerotic diseases. Overall, the inconsistency of association in the present and previous studies suggests that lipids only seem to play a minor part in the progression of AAA, and thus indirectly in the pathogenesis of AAA.

Table 1. Beta blockade, dietary and smoking habits as dichotomous variables tested concerning antibodies against oxidised low density lipoprotein, lipids, lipoprotein (A), annual change in ankle-brachial-blood pressure index (ABI), and annual expansion rate of small abdominal aortic aneurysm. Median values with 25% and 75% quartile values shown in parentheses. *p*-Values of Wilcoxon's rank sum test.

Variable	Y	N	S-Ab-oxLDL (units) Y/N	S-LDL (mmol/l) Y/N	S-HDL (mmol/l) Y/N	S-total cholesterol (mmol/l) Y/N	S-Lp(A) mmol/l Y/N	S-triglycerides (mmol/l) Y/N	ABI-change per year Y/N	Expansion rate (mm/year) Y/N
Beta-blockage	25	112	(17) 25 (39) (18) 21 (28) <i>p</i> =0.22	(0.9) 1.4 (1.6) (0.9) 1.1 (1.4) <i>p</i> =0.22	(0.9) 1.1 (1.2) (0.9) 1.1 (1.2) <i>p</i> =0.93	(5.8) 6.7 (7.1) (5.4) 6.0 (6.9) <i>p</i> =0.13	(2.5) 5.8 (31) (3.9) 10.3 (50) <i>p</i> =0.97	(2.3) 2.6 (3.1) (1.9) 2.4 (2.9) <i>p</i> =0.15	(1.0) 0.0 (-1.0) (1.0) -2.0 (-6.0) <i>p</i> =0.01	(1.0) 1.6 (2.7) (1.0) 2.5 (3.9) <i>p</i> =0.01
Regular eating habits	115	15	(18) 21 (30) (16) 21 (25) <i>p</i> =0.33	(0.9) 1.1 (1.4) (0.7) 1.0 (1.3) <i>p</i> =0.24	(0.9) 1.2 (1.4) (0.9) 1.1 (1.3) <i>p</i> =0.29	(5.5) 6.1 (7.0) (5.0) 6.0 (6.9) <i>p</i> =0.70	(3.0) 9.8 (29) (4.7) 8.5 (13) <i>p</i> =0.73	(1.9) 2.4 (2.9) (2.4) 2.6 (3.4) <i>p</i> =0.24	(1.0) -1.0 (-5.0) (1.5) 0.5 (-4.0) <i>p</i> =0.22	(1.0) 1.9 (3.2) (1.6) 2.8 (4.6) <i>p</i> =0.09
Tablet supply of vitamins	64	67	(18) 21 (28) (17) 22 (30) <i>p</i> =0.85	(0.9) 1.1 (1.4) (0.9) 1.1 (1.4) <i>p</i> =0.95	(1.0) 1.2 (1.3) (0.9) 1.2 (1.4) <i>p</i> =0.68	(5.4) 6.0 (7.0) (5.7) 6.2 (6.6) <i>p</i> =0.97	(3.7) 8.0 (29) (3.6) 10.5 (28) <i>p</i> =0.63	(2.0) 2.5 (3.0) (1.8) 2.4 (3.0) <i>p</i> =0.55	(1.0) -1.0 (-5.0) (1.0) -2.0 (-5.0) <i>p</i> =0.88	(1.1) 2.0 (3.1) (0.9) 1.9 (3.7) <i>p</i> =0.58
Tablet supply of Q10	12	110	(20) 21 (23) (17) 21 (28) <i>p</i> =0.86	(1.0) 1.2 (1.3) (0.9) 1.1 (1.5) <i>p</i> =0.90	(0.9) 1.2 (1.4) (0.9) 1.2 (1.3) <i>p</i> =0.93	(5.7) 6.1 (6.7) (5.5) 6.1 (7.0) <i>p</i> =0.95	(0.0) 2.5 (21) (4.1) 10.5 (29) <i>p</i> =0.04	(1.9) 2.4 (2.9) (1.9) 2.4 (2.9) <i>p</i> =0.89	(1.0) 0.5 (-1.0) (1.0) -2.0 (-6.0) <i>p</i> =0.04	(0.9) 2.0 (2.4) (1.0) 2.5 (3.9) <i>p</i> =0.19
Smoking	82	52	(17) 21 (29) (18) 22 (28) <i>p</i> =0.79	(0.8) 1.1 (1.4) (1.0) 1.1 (1.5) <i>p</i> =0.33	(0.9) 1.2 (1.3) (1.0) 1.2 (1.4) <i>p</i> =0.87	(5.5) 6.0 (6.5) (5.4) 6.3 (7.1) <i>p</i> =0.26	(2.8) 9.9 (28) (4.1) 10.1 (32) <i>p</i> =0.66	(1.9) 2.3 (2.8) (2.0) 2.5 (3.2) <i>p</i> =0.36	(1.0) -1.0 (-6.0) (0.0) -2.0 (-4.0) <i>p</i> =0.42	(1.0) 2.6 (4.1) (0.9) 1.7 (2.8) <i>p</i> =0.01*

\*: *p*<0.05 (*p*-value of multivariate regression analysis adjusting for smoking). Only performed on univariate significant tests.

Table 2. Spearman's non parametric correlation coefficients of selected continuous variables focusing on aneurysmal expansion rate and rate of decreasing ankle blood pressure index (ABI) as dependent variables.

	AAA-size (mm)	Initial ABI (%)	ABI- change (%/year)	S-LDL (mmol/l)	S-HDL (mmol/l)	S-tri- glycerides (mmol/l)	S-total cholesterol (mmol/l)	Ab-oxLDL (units/l)	Lp(a) (mmol/l)	Body mass index (kg/w <sup>2</sup> )
Initial AAA-size (mm)	1.00	0.11 (0.21)	0.11 (0.73)	0.013 (0.99)	0.10 (0.32)	0.01 (0.92)	0.01 (0.90)	-0.16 (0.08)	-0.04 (0.74)	0.03 (0.75)
AAA-expansion (mm/year)	0.30 (0.01)*	0.06 (0.50)	0.11 (0.20)	0.031 (0.97)	0.01 (0.91)	0.21* (0.04)	0.17 (0.14)	-0.15 (0.09)	0.11 (0.33)	-0.01 (0.94)
Initial ABI	0.11 (0.21)	1.00	0.33 (0.01)*	-0.01 (0.98)	0.01 (0.89)	-0.07 (0.51)	-0.09 (0.43)	-0.12 (0.20)	-0.18 (0.06)	0.09 (0.30)
ABI-change (per cent/year)	0.11 (0.73)	0.33 (0.01)*	1.00	0.01 (0.98)	0.01 (0.91)	-0.11 (0.31)	-0.04 (0.73)	-0.15 (0.11)	0.26 (0.02)	0.04 (0.64)

ABI: ankle brachial systolic blood pressure index.

HDL: High density lipoprotein.

LDL: Low density lipoprotein.

Ab-oxLDL: antibodies against oxidised LDL.

Lp(a): lipoprotein A.

\*:  $p < 0.05$  after multiple linear regression analysis adjusting for smoking. (Only performed on univariate significant tests, and  $\log(\text{Lp(a)})$  concerning  $\text{Lp(a)}$ ).

The concentration of antibodies against malondialdehyde-modified LDL has been shown to be significantly elevated in patients with AMI and carotid-stenosis, but not in angina pectoris,<sup>37</sup> and positively correlated to the progression of carotid stenosis. This correlation was increased by presence of hypertriglyceridaemia.<sup>5</sup> However, the level of ab-oxLDL could not be associated to aneurysmal progression. This is hardly due to a large variation of measurement because only one scanner and two observers were used. Validation of our measurements have earlier been reported,<sup>13</sup> 95% of their measurements are within 2 mm difference.<sup>14</sup> Furthermore, the mean average expansion rate was 2.6 mm/year, and the average observation time was 2.5 years. Consequently, a clinical significant correlation to expansion must be expected to be detectable. Also, Ab-oxLDL were not associated with a faster decrease in ABI. However, the ABI changes were relatively minor compared with expansion rates, and the variation of the ABI measurements was larger (15% of the measurements are measured within 15% difference).

The negative results concerning ab-oxLDL cannot be taken as a sign of minor pathogenetic importance of oxidised LDL because plasma oxidised LDL seems to increase the transposition through the arterial endothelium and wall, and to increase the degradation of the wall.<sup>38</sup> Consequently, oxidised LDL could play a part in the aneurysmal and atherosclerotic process. The negative results concerning Ab-oxLDL may rather question whether the antibodies will reflect the degree of this transposition because oxidized LDL seems only present in the circulation for minutes.<sup>38</sup>

Finally, lipoprotein (A) levels have earlier been correlated to the incidence of intermittent claudication.<sup>1</sup> By adding objective and prospective measurements, our results supports this, while the lack of any correlation to aneurysmal expansion indicates partly different pathogenetic factors between atherosclerosis and aneurysmal degradation.

### Conclusion

If anything, lipids seem to play only a minor role in the progression of AAA, and pharmacological intervention to lower the serum levels can only be justified in order to prevent atherosclerotic events. In contrast, smoking cessation may inhibit aneurysmal expansion. Antibodies against oxidised LDL seem not to be predictive of aneurysmal or atherosclerotic progression,

and can at this point not be recommended for monitoring or decision-making concerning antioxidant therapy. However, the role of oxidised low density lipoprotein remains unsolved.

### Acknowledgements

The Danish Heart Foundation, The Foundation of Asta and Rosa Jensen, and Viborg County are thanked for financial support. The nurse of the trial, Anette Sahlholdt, Dr Ulrik Gerdes, University of Aarhus, and Dr Per Jæger, Skive Sygehus, are thanked for excellent and committed practical and administrative support.

### References

- 1 BOSTOM AG, GAGNONN DR, CUPPLES LA *et al.* A prospective investigation of elevated lipoprotein a detected by electrophoresis and cardiovascular disease in women. The Framingham Heart Study. *Circulation* 1994; **90**: 1688–1695.
- 2 HOVOET P, PEREZ G, ZHAO Z *et al.* Malondialdehyde-modified low-density-lipoproteins in patients with atherosclerotic disease. *J Clin Invest* 1995; **95**: 2611–2619.
- 3 SALONEN JT, HERTTUALA SY, YAMAMOTO R *et al.* Autoantibodies against oxidised LDL and progression of carotid atherosclerosis. *Lancet* 1992; **339**: 883–887.
- 4 GOODE GK, MILLER JP, HEAGERTY AM. Hyperlipidaemia, hypertension, and coronary heart disease (Review). *Lancet* 1995; **345**: 362–364.
- 5 REGNSTRÖM J, NILSSON J, TORNVALL P, LANDOU C, HAMSTEN A. Susceptibility to low-density lipoprotein oxidation and coronary atherosclerosis in man. *Lancet* 1992; **339**: 1183–1186.
- 6 LINDHOLT JS. Considerations and experiences of screening for abdominal aortic aneurysms (Ph.D.thesis). ISBN: 87-7749-185-8. Copenhagen: FADL's Forlag, 1998.
- 7 VAARALA O, ALFTHAN G, JAUHAINEN M *et al.* Cross reaction between antibodies to oxidised low-density lipoprotein and to cardiolipin in systemic lupus erythematosus. *Lancet* 1993; **341**: 923–925.
- 8 PALINSKI W, YLÄ-HERTTUALA S, ROSENFELD ME *et al.* Antisera and monoclonal antibodies specific for epitopes generated during oxidative modification of low density lipoprotein. *Atherosclerosis* 1990; **10**: 325–335.
- 9 KLAUSEN IC, NIELSEN FE, HEGEDÜS L *et al.* Treatment of hypothyroidism reduces low-density-lipoproteins but not lipoprotein(a). *Metabolism* 1992; **41**: 911–914.
- 10 FRIEDEWALD WT, LEVY RI, FREDRICKSON DS. Estimation of the correlation of low-density lipoprotein cholesterol in plasma without use of the preparative ultracentrifuge. *Clin Chemistry* 1972; **18**: 499–502.
- 11 LINDHOLT JS, JUUL S, FASTING H, HENNEBERG EW. Is screening for abdominal aortic aneurysm acceptable for the population? Selection and recruitment for hospitalbased screening for abdominal aortic aneurysm. *J Public Health Med* 1998; **20**: 211–217.
- 12 LINDHOLT JS, HENNEBERG EW, FASTING H, JUUL S. High-risk or mass screening for abdominal aortic aneurysms. *Br J Surg* 1997; **84**: 40–42.
- 13 LINDHOLT JS, VAMMEN S, JUUL S, FASTING H, HENNEBERG EW. The validity of ultrasonographic scanning as screening method for abdominal aortic aneurysm. *Eur J Vasc Endovasc Surg* 1999; **17**: 472–475.
- 14 LINDHOLT JS, VAMMEN S, LIND I, HENNEBERG EW, FASTING H. Progression of lower limb atherosclerosis is associated with antibodies against *Chlamydia pneumoniae*. *Eur J Vasc Endovasc Surg* 1999; **18**: 527–529.

- 15 BENGSSON H, BERGQVIST D, STERNBY NH. Increasing prevalence of abdominal aortic aneurysms. A necropsy study. *Eur J Surg* 1992; **158**: 19–23.
- 16 STERPETTI AV, SCHULTZ RD, FELDHAUS RJ et al. Abdominal aortic aneurysms in elderly patients. Selective management based on clinical status and aneurysmal expansion rate. *Am J Surg* 1985; **150**: 772–776.
- 17 BENGSSON H, NILSSON P, BERGQVIST D. Natural history of abdominal aortic aneurysm detected by screening. *Br J Surg* 1993; **80**: 718–720.
- 18 COLLIN J, HEATHER B, WALTON J. Growth rates of subclinical abdominal aortic aneurysms – implications for review and re-screening Programmes. *Eur J Vasc Surg* 1991; **5**: 141–144.
- 19 BENGSSON H, BERGQVIST D, EKBERG Ö, RANSTAM J. Expansion pattern and risk of rupture of abdominal aortic aneurysms that were not operated on. *Eur J Surg* 1993; **159**: 461–467.
- 20 STERPETTI AV, SCHULTZ RD, FELDHAUS RJ, CHENG SE, PEETZ DJ. Factors influencing enlargement of small abdominal aortic aneurysms. *J Surg Research* 1987; **43**: 211–219.
- 21 CRONENWETT J. Variables that affect the expansion rate and outcome of small abdominal aortic aneurysms. *J Vasc Surg* 1990; **11**: 261–269.
- 22 PLEUMEEKERS HJCM, HOES AW, VAN DER DOES E et al. Aneurysms of the abdominal aorta in older adults. The Rotterdam study. *Am J Epidemiol* 1995; **142**: 1291–1299.
- 23 FRANKS PJ, EDWARDS RJ, GREENHAULGH RM, POWELL JT. Risk factors for abdominal aortic aneurysms in smokers. *Eur J Vasc Endovasc Surg* 1996; **11**: 487–492.
- 24 KROHN CD, KULLMANN G, KVERNEBO K, ROSEN L, KROESE A. Ultrasonographic screening for abdominal aortic aneurysm. *Eur J Surg* 1992; **158**: 527–530.
- 25 LOUWRENS HD, ADAMSON J, POWELL JT, GREENHAULGH RM. Risk factors for atherosclerosis in men with stenosing or aneurysmal disease of the abdominal aorta. *Int Angio* 1993; **12**: 21–24.
- 26 SMITH FCT, GRIMSHAW GM, PATERSON IS, SHEARMAN CP, HAMER JD. Ultrasonographic screening for abdominal aortic aneurysm in an urban community. *Br J Surg* 1993; **80**: 1406–1409.
- 27 KELLY TJ, HEATHER BP. General practice-based population screening for abdominal aortic aneurysm. *Br J Surg* 1989; **76**: 479–480.
- 28 AUERBACH O, GARFINKEL L. Atherosclerosis and aneurysm of aorta in relation to smoking habits and age. *Chest* 1980; **78**: 805–809.
- 29 MACSWEENEY ST, ELLIS M, WORRELL PC, GREENHAULGH RM, POWELL JT. Smoking and growth rate of small abdominal aortic aneurysm. *Lancet* 1994; **344**: 651–652.
- 30 LEACH SD. Effect of B-adrenergic blockade on the Growth rate of Abdominal Aortic Aneurysms. *Arch Surg* 1988; **123**: 606–609.
- 31 GADOWSKI GR, PILCHER DB, RICCI MA. Abdominal aortic aneurysm expansion rate: effect of size and beta-adrenergic blockade. *J Vasc Surg* 1994; **19**: 727–731.
- 32 BOUCEK RJ. Modulation by propranolol of lysyl cross links in aortic elastin and collagen of the aneurysm-prone turkey. *Biochemical Pharmacology* 1983; **32**: 275–280.
- 33 LINDHOLT JS, FASTING H, JUUL S, HENNEBERG EW. Impaired results of a randomised double blinded trial of propranolol versus placebo on the expansion rate of small abdominal aortic aneurysms. *Int Angio* 1999; **18**: 52–57.
- 34 KRUPSKI WC. Utility of computer tomography for surveillance of small abdominal aortic aneurysms. *Arch Surg* 1990; **125**: 1345–1450.
- 35 NORRGAARD Ö. Familial Aortic Aneurysms: Serum concentrations of triglyceride, cholesterol, HDL-cholesterol, and (VLDL + LDL)-cholesterol. *Br J Surg* 1985; **72**: 113–116.
- 36 BENGSSON H, EKBERG Ö, ASPELIN P, KÄLLERÖ S, BERGQVIST D. Ultrasound screening of the abdominal aorta in patients with intermittent claudication. *Eur J Vasc Surg* 1989; **3**: 497–502.
- 37 JONES HJS, IBRAHIM AEK, HOSKINS C, DERODRA JK. Aortic screening in general practice. *Br J Surg* 1996; **348**: 1320.
- 38 NIELSEN LB. Atherogenicity of lipoprotein (a) and oxidized low density lipoprotein (Thesis). Copenhagen: University of Copenhagen, 1999.

Accepted 11 November 2000